


BIOPIROX

10 mg/ml

cutaneous spray, solution

Piroctone olamine

100 ml



bioveta



Biopirox is a clear to slightly opalescent cutaneous solution containing in 1 ml: **Active substance:** Piroctone olamine – 10 mg • **INDICATIONS:** Treatment of mycotic skin diseases caused by dermatophyte fungi in dogs, cats, furry and small animals. • **CONTRAINDICATIONS:** None. • **ADVERSE REACTIONS:** Possible allergic reactions in particularly susceptible individuals. A 10-fold increase in dose causes a local reaction, formation of erythema and alopecia at the injection site. These changes resolve 5–7 days later - erythema subsides, the skin becomes smooth and animal's hair begins to recover. • Treated animal manifests no general reaction after administration of the product. • **TARGET SPECIES:** Dog, cat, non-food furry and small non-food animals. • **DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:** The product should be applied to affected sites by spraying from a distance of 10 cm, at least 4 times in 2 to 4-day intervals until disappearance of clinical symptoms. • **Method of administration** –dermal, by spraying. • **ADVICE ON CORRECT ADMINISTRATION:** The highest daily dose and the maximum dose for the treatment: It is recommended to apply the product repeatedly not earlier than 2-4 days after the first application and to one quarter of the body as a maximum. • **WITHDRAWAL PERIOD:** Not intended for food animals. • **SPECIAL STORAGE PRECAUTIONS:** Keep out of the reach of children. Store below 25 °C. Protect from frost. • Do not use after the expiry date which is stated on the packaging. • **SPECIAL WARNING(S):** Avoid contact with skin, eyes and mucous membranes. After contact with skin or mucosa, wash immediately with plenty of water, take off all contaminated clothing which is in direct contact with skin. In case of contact with eyes, rinse with plenty of drinking water. In case of irritation seek medical attention. • Avoid inhalation of the product. Use of personal protective equipment including neoprene gloves is recommended for repeated applications (e.g. in breeding facilities, shelters). • In case of accidental ingestion drink plenty of water and seek medical advice immediately and show the package leaflet or label to a practitioner. • Do not smoke, eat or drink during the application. • There are no known interactions with other medicines. Generally, it is not recommended to initiate active immunisation against mycoses concomitantly with antifungal therapy. • The principle of active immunisation consists in injection of live attenuated or inactivated strains of fungal agent and concomitant application of BIOPIROX spray can reduce the immune response to vaccination. • **SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS:** Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED: May 2010

bioveta

