

DEXACAL 40 HAMPTON

Treatment of inflammation | steroid anti-inflammatory

Veterinary use. To be sold with a veterinary prescription.

Always consult with your Veterinary Doctor.

NET CONTENT: 10 and 50 mL



Calastremé
PRODUCTOS VETERINARIOS

MADE IN ARGENTINA

FORMULA:

Dexamethasone 21 Phosphate 4 mg
Sterile vehicle q.s.....1 mL

ACTION:

Dexacal 40 Hampton is recommended in those clinical processes that require aggressive and continuous corticosteroid therapy. Local and general tolerance is excellent.

DOSES AND METHOD OF ADMINISTRATION:

By intramuscular or intravenous injection.

Equines: 1,25 to 2,5 mL per day (0,05 mg per kg bodyweight).

Bovines: 2,5 to 10 mL per day (0,02 mg per kg bodyweight). Labor Induction in cows: 10 mL

Porcines: 0,5 to 2 mL per day (0,01 mg per kg bodyweight).

Canines and Felines: 0,05 to 2 mL per day (0,1 mg per kg bodyweight).

SPECIES:

Equines, Bovines, Porcines, Canines and Felines.

INDICATIONS FOR USE:

In all species: anti-inflammatory antiallergic corticosteroid and increase tolerance to toxins in case of infections.

Equines : arthritis, tenosynovitis, lumbago, bursitis, chronic obstructive pulmonary disease (COPD), pododermatitis, azoturia.

Bovines: Ketosis, puerperal paresis, pre and postpartum paraplegia, tetanias, arthritis, bursitis, mastitis, metritis
Porcines: Agalactia, arthritis, eclampsia, intoxications, dermatosis disease, shock prevention, mastitis-metritis-agalactia.

Canines y felines: Eczemas, dermatitis, non-septic arthritis, inflammatory and / or allergic conditions, intervertebral disk syndrome.

WARNING:

Shake before use. Do not exceed the recommended dose.
Do not use in gastrointestinal ulcers, Cushing's disease,

osteoporotic processes, diabetes mellitus, active tuberculosis and viral infections.

It should not be administered to pregnant females during the last third of gestation

PRECAUTIONS:

DEXACAL 40 HAMPTON should be used with caution in patients with infections, kidney failure, corneal ulcers, hypotension, osteoporosis, diabetes mellitus and congestive heart failure.

Not recommended in pregnant females.

Do not stop treatment abruptly after prolonged treatment, as this may cause adrenocortical insufficiency.

It is advisable to reduce the dose gradually.

PRESENTATION:

Vials 10 and 50 mL

NOTE: Since the product will be used in circumstances beyond the control of the supplier, the latter is not to be held responsible for the consequences arising from the improper use thereof.

Keep out of reach of children and pets.

Storage conditions: Store between 15° to 25°C, protected from light.

Used containers must be discarded in accordance with current local legislation.

In the event of accidental poisoning attend the nearest poison control center.

Withdrawal time :

For meat : 3 days

For milk : 1 day



SENASA Cert. N° 07-212
Estab. Elab. N° 8533



PROTECT
THE ENVIRONMENT



ARGENTINE PRODUCT



Calastremé
PRODUCTOS VETERINARIOS

Produced and distributed by : Establecimiento Calastremé S.A. La Paz 2150 B1640CYB
Martínez, Buenos Aires, Argentina. Ph.: (011) 4580-9999 e-mail: info@calastreme.com.ar
www.calastreme.com.ar