

DIFEBUZOL

ANALGESIC ANTI-INFLAMMATORY

Presentation: 100 mL bottle

Formula: Every 100 ml contains: Phenylbutazone 20 g; Excipient q.s. 100 ml

Indications:

Analgesic, anti-inflammatory and antipyretic, indicated to relieve inflammatory processes and pain caused by lesions of the musculoskeletal system.

Myalgia, tendon injuries, post-traumatic pain. Adjuvant treatment of colic (to reduce endotoxin effects).

Adjuvant treatment of laminitis.

Administration and dosage:

Injectable (slow intravenous administration only).

In general, 3 to 6 mg/kg every 12 hours, not exceeding 8.8 mg/kg per day.

Adjuvant treatment of colic (to reduce endotoxin effects): 2.2 mg/Kg.

Adjuvant treatment of laminitis 4.4 mg/Kg.

For an adult horse, 10 ml (2 g of phenylbutazone) are normally applied every 24 hours.

For severe cases, the dose can be divided in two to improve the plasma profile. Do not exceed 5 days of treatment.



Contraindications and precautions:

Contraindicated in horses with a spinal or hematological history and previous gastrointestinal ulcers.

Do not use it in animals intended for human consumption. Do not use it in animals with hypersensitivity disorders.

Administer with caution in foals and pony breeds due to the higher incidence of hypoproteinemia and ulcers, and in patients with a renal and cardiac history.

Target species: Equines not intended for human consumption.

To be sold with a veterinary prescription

Keep out of the reach of children and pets.

National Poison Control Center: 0800-333-0160

Storage: Between 15° and 25°C and sheltered from light.

Used containers must be discarded according to current local legislation.



Calastremé