

FORMULATION:

Each g contains:

Neomycin sulfate4.58 mg
Tetracycline Hydrochloride 4.2 mg
Prednisolone Acetate 0.5 mg
Excipient (Vitamin E) q.s. 1 g

**CONSULT A
VETERINARY DOCTOR**

**AVAILABLE BY
PRESCRIPTION ONLY**

NET CONTENT:
8, 10, 12, 25, 50, 100, 120 y
150 g

NEOSIN-T

**REGISTRATION NUMBER
Q-7972-191**

VETERINARY USE

Ointment

Tetracycline, neomycin and
prednisolone

Broad-spectrum bactericide and
anti-inflammatory agent



MADE IN MEXICO BY:
PET'S PHARMA DE MÉXICO
S.A. DE C.V.
Av. Sor Juana Inés de la Cruz
N° 580,
Col. Benito Juárez,
Nezahualcóyotl
Estado de México, México.,
'P.C. 57000
Ph.: 55 57 43 38 39
www.petspharma.com.mx
ventas@petspharma.com.mx

INDICATIONS:

Indicated for the treatment of simple or infected wounds, sores, burns, pustular dermatitis and eczema. The combination of Neomycin and Tetracycline has broad-spectrum bactericidal properties, it attacks the main pathogenic microorganisms involved in infections and skin wounds. Prednisolone Acetate facilitates a rapid evolution of wounds due to its anti-inflammatory action. It is efficient against Gram positive bacterias: *Corynebacterium* spp., *Listeria monocytogenes*, *Clostridium* spp., *Bacillus anthracis*, *Staphylococcus* spp.; Gram negative bacterias: *Escherichia coli*, *Brucella* spp., *Plesiomonas* spp., *Aeromonas* spp., *Neisseria* spp., *Klebsiella* spp., *Proteus* spp., *Pseudomonas aeruginosa*, *Salmonella* spp., *Enterobacter* spp., *Serratia* spp., *Shigella* spp., *Bacteroides fragilis*.

SPECIES

Bovine, Equine, Ovine, Caprine, Swine, Canine and Feline

DOSE:

3 to 4 times for day at Veterinary Doctor's discretion.

INSTRUCTIONS FOR USE:

To Apply a considerable amount over the affected area.

ROUTE OF ADMINISTRATION:

Topical.

WARNINGS AND/OR RESTRICTIONS:

External use, do not ingest. In case of accidental ingestion, show the product label to the doctor. It is not considered a toxic product, however, depending on the amount of product ingested, it can cause diarrhea and/or vomiting. Avoid contact with the eyes. Discontinue immediately in case of adverse reactions. Keep in a cool, dry place, in a range within 15-30 °C and protected from direct sunlight. Keep out of reach of children and pets. Any unused veterinary drug product, residues derived from it and empty containers must be disposed safely and in accordance with local regulations, avoiding contaminating water sources. **Product for exclusive use in Veterinary Medicine.**