

Protecto Entero Pectin[®]

 Laboratorio
FUNDACION
CIENCIA Y TECNOLOGÍA FARMACEUTICA

STORE BETWEEN 15 AND 25 °
VETERINARY USE.
FREE SALE.

ANTIDIARRHEAL
Con. Neto 100 ml

COMPOSITION:

Bismuth Subnitrate_ 3gr / Aluminum Silicate_ 30gr / Pectina_0.5gr / Dimethicone_1gr / Excipient c.s.p. 100gr

THERAPEUTIC ACTION:

antidiarrheal, backmolytic, protector of the mucosa, antiseptic, adsorbent of toxins, and carminative.

INDICATIONS:

Diarrhea of any etiology, food poisoning enteritis, gastroenteritis, gastritis, and meteorism.

ADMINISTRATION: Oral.

DOSAGE:

0.5 to 2 ml / kgpv in equines, bovines, sheep, goats, pigs, and felines; or criteria of the acting veterinarian. It can be repeated every 12 hours. The eventual presence of lumps does not alter the quality or efficacy of the product. Shake before using.

SIDE EFFECTS: Protecto Entero Pectin does not have side effects.

OBSERVATIONS:

It's not recommended to be mixed with milk and/or milk substitute, but it is advisable to give it half an hour before them. Oral rehydration salts shouldn't be mixed with the product but given after or before the treatment.

