

ACEPROMAZINA HAMPTON

Injectable ataraxic, tranquilizing.



Calastremé
PRODUCTOS VETERINARIOS

Veterinary use. To be sold with a field veterinary prescription.

NET CONTENT: 10 and 50 mL

MADE IN ARGENTINA

FORMULA:

Acepromazine Maleate 1 g.
Excipients q.s. 100 mL

INDICATIONS:

Generally indicated as preanesthetic medication, being a powerful neuroleptic agent.
As an adjuvant in the control of unruly animals and/or of those presenting trauma that hinders reviewing them at the time of clinical examination (e.g. fractures, burns, etc.). It can also be used prior to transporting and/or loading. It is useful when used together with other local or general anesthetics for various procedures or treatments (castration, neurectomies, removal of skin tumors, eye surgeries, problems related to shoeing, etc.). It can be used in equines as adjuvant treatment of laminitis (developmental phase) and in cases of colic, to restore bowel transit.

DOSES AND ROUTES OF ADMINISTRATION:

Each ml of product contains 10 mg of Acepromazine.
The doses to be administered according to species are:
Canines: 0.05-0.1 mg/kg, slow IV, IM, SC (Do not exceed a total dose of 3 mg). Equivalent to 0.05-0.1 ml of Acepromazine Hampton per 10 kg of weight. (Do not exceed a total of 0.3 ml).
Felines: 0.05-0.1 mg/kg slow IV, IM, SC (Do not exceed a total dose of 1 mg). Equivalent to 0.025-0.05 ml of Acepromazine Hampton per 5 kg of weight. (Do not exceed a total of 0.1 ml)
Equines: 0.01-0.05 mg/kg slow IV or IM 0.044-0.088 mg/kg SC. Equivalent to 0.4-2 ml IV or IM or 1.75-3.5 ml SC per 400 kg weight.

CONTRAINDICATIONS AND USAGE LIMITATIONS:

Acetylpromazine has an adrenolytic action by blocking alpha receptors and thus reverses the pressor effect of adrenaline, which acts only on beta receptors. Therefore, adrenaline should not be used as a pressor amine, but rather norepinephrine or phenylephrine, which only act on alpha receptors.
It is advisable to use it with caution in animals with liver disease, since the ACEPROMAZINA HAMPTON component

is broken down in the liver.

It can cause minor cardiac arrhythmias. It is advisable to use with caution in animals with history of cardiac disorders. In animals with heart disease, it can increase its intensity, so it is advised to use it with caution as well as in senile animals. Apply slowly intravenously. In epileptic animals, it can cause seizures. Its use is not recommended.

Like all phenothiazine derivatives, it potentiates the sensitivity of organophosphorus compounds and that of procaine hydrochloride.
For this reason, their simultaneous use should be avoided. Occasionally, adult stallions may suffer from irreversible prolapse of the penis, which must be reset manually.

NOTE: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.

Keep out of the reach of children and pets.
Store between 15° and 25°C and sheltered from light.
Used containers must be discarded in accordance with current local legislation.
Discard leftover product once the container has been opened.
In the event of accidental poisoning attend the nearest poison control center.

Technical Director: Dr. Carlos N. Nuñez. Veterinary Doctor.
Medical Registration P.B.A 1350



SENASA Cert. N° 80-287
Estab. Elab. N° 8533



PROTECT
THE ENVIRONMENT



ARGENTINE PRODUCT