



Amprocox WS

Powder for oral administration

Composition:

Contains per gram powder:

Amprolium hydrochloride.....	200 mg.
Sulfaquinoxaline	150 mg.
Vitamin A, retinol acetate	15 000 IU.
Vitamin K ₃	5 mg.
Excipients ad	1 g.

Description:

Amprolium is an anticoccidial with activity against *Eimeria* spp., especially *Eimeria tenella* and *Eimeria necatrix*. Sulfaquinoxaline is a chemotherapeutic with a bacteriostatic activity against many Gram-positive and Gram-negative bacteria and has also coccidiostatic activity against *Eimeria* spp. Vitamins are essential for the proper operation of several physiological functions.

Indications:

Coccidiosis in calves, sheep, goats and poultry.

Contra-indications:

Administration to animals with impaired hepatic and/or renal functions.
Hypersensitivity to amprolium and/or sulfaquinoxaline.

Side effects:

At high dosages in laying hens egg-drop and in broilers growth inhibition and polyneuritis can occur. Other side effects may include crystalluria, anaemia, leucopenia and thrombocytopenia.

Dosage:

For oral administration:

Calves, sheep and goats:

- Preventive : 1 g per 50 - 100 kg body weight through drinking water or milk for 21 days.
- Curative : 5 g per 25 - 50 kg body weight through drinking water or milk for 5 days.

Poultry : 20 g per 20 - 40 litres of drinking water for 5 - 7 days.

Note: for pre-ruminant calves, lambs and kids only. Do not administer to hens producing eggs for human consumption.

Withdrawal times:

- For meat:

- Calves, sheep and goats : 14 days.
- Poultry : 14 days.

Packaging:

Sachet of 100 g and jar of 1000 g.

