




PACKAGE LEAFLET FOR:

Ancesol 10 mg/ml solution for injection for cattle

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT

Marketing authorisation holder and manufacturer responsible for batch release:

VetViva Richter GmbH, Durisolstrasse 14, 4600 Wels, Austria

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Ancesol 10 mg/ml solution for injection for cattle

Chlorphenamine maleate

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

1 ml contains:

Active substance:

Chlorphenamine maleate	10 mg
(equivalent to 7.03 mg chlorphenamine)	

Excipients:

Methyl parahydroxybenzoate (E218)	1.00 mg
Propyl parahydroxybenzoate	0.20 mg

Clear, colourless to almost colourless solution

4. INDICATION(S)

For the symptomatic treatment of conditions associated with histamine release.

5. CONTRAINDICATIONS

Do not use in case of hypersensitivity to the active substance or to any of the excipients.

6. ADVERSE REACTIONS

Chlorphenamine has a weak sedative effect.

If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.

7. TARGET SPECIES

Cattle

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

For intramuscular or slow intravenous use, see also section "9. Advice on correct administration".

Adult animals:

0.5 mg Chlorphenamine maleate/kg bodyweight (5 ml/100 kg bodyweight), once a day for three consecutive days.

Calves:

1 mg Chlorphenamine maleate/kg bodyweight (10 ml/100 kg bodyweight), once a day for three consecutive days.

9. ADVICE ON CORRECT ADMINISTRATION

Although intravenous administration has an immediate therapeutic effect, it can have excitatory effects on the CNS. Consequently, administer slowly and interrupt administration for a few minutes if necessary, when using this route.

Do not administer via the subcutaneous route.

10. WITHDRAWAL PERIOD

Meat and offal: 24 hours

Milk: 12 hours

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the label and carton after EXP. The expiry date refers to the last day of that month.

Shelf-life after first opening the container: 28 days

Do not store above 30 °C.

12. SPECIAL WARNING(S)

Special warnings for each target species

None