

ARTROCAL PLUS HAMPTON

PREVENTIVE, THERAPEUTIC, RESTORATIVE SUPPLEMENT | ORAL POWDER

Presentation: 500 and 1000g. Pot

Formula: Glucosamine hydrochloride 27.50 g.; Chondroitin sulfate(*) 10.00 g.; Manganese sulfate(*) 0.70 g.; Ascorbic acid 1.60 g.; Excipients q.s. 100 g.

Indications:

Especially indicated to treat joint disorders caused by training in sport horses. Osteoarticular restorer, chondrogenesis promoter. Arthrosis, non-infectious arthritis, osteoarthritis, hydrarthrosis and other degenerative arthropathies, chondropathies, synovitis, osteoarticular tendinitis, coadjuvant in fracture healing and post osteoarticular surgery. Osteochondrosis. Joint anti-inflammatory.

Dosage and method of administration:

The product is administered orally, on its own or mixed with a ration. If mixed with a ration, discard what is not consumed on the same day it is prepared. Do not re-use. The product comes with a graduated measuring spoon.

Loading dose:

In horses weighing less than 300 kg, administer 13 g in the morning and 13 g in the afternoon. In horses weighing between 300 to 600 kg, administer 20 g in the morning and 20 g in the afternoon. In horses weighing over 600 kg, administer 28 g in the morning and 28 g in the afternoon.

Maintenance dose:

In horses weighing less than 300 kg, administer 6.6 g a day.

In horses weighing between 300 to 600 kg, administer 6.6 to 13g a day.

In horses weighing over 600 kg, administer 13 g a day.

During this period, the dose may be increased up to the loading dose, depending on the animal's response.

Administer one dose in the morning and one in the afternoon.

Treatment duration:

Thirty days. In case of not obtaining a good response, the treatment may be extended for another 15 days.

Caution:

The product should not be administered to pregnant females because its innocuousness has not been proven.

The product should not be applied to animals presenting a history of hypersensitivity to any of the components.

Do not administer to horses intended for human consumption.

Keep out of the reach of children and pets. In the event of accidental poisoning attend the nearest poison control center.

Storage: Between 15° and 25° C and sheltered from light.

Used containers must be discarded according to current local legislation.



Calastremé