

Biocan M inj. ad us. vet.

Vaccine against *Microsporum canis* in dogs and cats

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE

Bioveta, a. s.
Komenského 212
683 23 Ivanovice na Hané
Czech Republic

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

Biocan M inj. ad us. vet.
Vaccine against *Microsporum canis* in dogs and cats

3. STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS

Composition – 1 ml (1 dose) contains:

Active substance:
Microsporum canis inact.
min. 500 000 vegetative forms

Excipients:

Algeldratum
Sodium chloride solution

4. INDICATION

From the 12th week of age for the prophylaxis and therapy of dermal mycosis in dogs and cats caused by the dermatophyte *Microsporum canis*.

5. CONTRAINDICATIONS

Generalized fever. Vaccination of pregnant bitches and cats is not recommended.

Further immunoprophylactic actions during one week before the first vaccination until the 14th day after the second (or third) vaccination (apart from Biocan vaccines administration).

6. ADVERSE REACTIONS

Adequate local reaction (of pea-size usually) may occur at the site of injection that subsides within 3 weeks. Hypersensitivity may occur in exceptional cases.

7. TARGET SPECIES

Dog, cat.

8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

1 ml regardless of age, weight or breed of the animal.

Dogs: strictly intramuscularly into the hind limb muscle.

Cats: subcutaneously into the area behind the blade bone or intramuscularly into the hind limb muscle.

Preferably vaccinate into the left and revaccinate into the right half of the body.

Primovaccinated animals should be revaccinated at least once at an interval of 14 – 21 days. In the case of therapeutic vaccination, apply another (third) vaccination dose in 18 – 24 days after revaccination, if necessary.

Cellular and partly humoral immunity is induced in immunized animals.

The immunity develops within 1 month after revaccination and persists for at least 1 year. Annual revaccinations are recommended to maintain permanent immunity.

9. ADVICE ON CORRECT ADMINISTRATION

Shake the content of the vial before use.

10. WITHDRAWAL PERIOD

Not designed for food-producing animals.

11. SPECIAL STORAGE PRECAUTIONS

Store in a dry and dark place at 2 to 8°C. Do not freeze.

12. SPECIAL WARNING

Do not use the product after the expiry date indicated on the cover.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS

Dispose of containers and any residues of the product in accordance with applicable national regulations.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED

July 2006

15. OTHER INFORMATION

Pharmacotherapeutic group:

Veterinary immunopreparation.

Dosage form:

Injection

Interactions:

Always apply Biocan M separately (never use it as a diluent for lyophilized vaccines or mix with liquid Biocan vaccines).

Any other Biocan vaccine (Puppy, P, DP, DH, DHP, DHPPI, L, R, LR, C) can be administered simultaneously with Biocan M into a different site (preferably on the other side).

Package size:

A/ Plastic box with a lid with 20 pits:

20 × 1 ml of Biocan M vaccine

10 × 1 ml of Biocan M vaccine

2 × 1 ml of Biocan M vaccine

B/ Plastic box with a lid with 100 pits:

100 × 1 ml of Biocan M vaccine

50 × 1 ml of Biocan M vaccine

bioveta