



Division: Injection

®

BRIT GENT-15

COMPOSITION

Each 1 mL contains:

Gentamicin 150 mg.

INDICATIONS

Gastrointestinal & respiratory infections caused by gentamicin sensitive bacteria, like *E. coli*, *Klebsiella*, *Pasteurella* and *Salmonella* spp. in livestock. Gentamicin exerts pronounced bactericidal activity against several, often multi-resistant, Gram-negative pathogens.

CONTRA-INDICATIONS

Do not administer to animals hypersensitive to amino-glycoside antibiotics. Do not administer to animals with an impaired renal function or with vestibular, ear or visual dysfunctions. Do not use in animals in lactation if milk is intended for human consumption.

SIDE EFFECTS

Kidney damage and/or toxicity, with the vestibular function most often being damaged, when given at high doses and during long times. Respiratory paralysis due to neuromuscular blockade (at very high doses) may occur.

DOSAGE

By intramuscular, subcutaneous or slow intravenous injection

- Horses, Cattles, Buffaloes: 1mL per 20 - 40kg of body weight for 3 days.
- Foals, Calves, Swine: 1 - 2mL per 50kg of body weight for 3 days.
- Piglets, Dogs, Cats, Poultry: 0.5mL per 5kg of body weight

By intra-uterine application

- Horses (only during estrus, Acute metritis, Chronic metritis): 6mL
- Cattle: 2mL

Recommended twice on the first day, half a dose every 12 hours from second day.

INCOMPATIBILITY WITH OTHER DRUGS

Do not combine with potential nephrotoxic or muscle paralyzing medicines.

WITHDRAWAL TIME

Meat: Cattle, sheep, goats: 90 days.

Pigs: 40 days.

DOSAGE

For intramuscular or intravenous injection.

Cattle:

30-40 ml/adult animal/day; and
10-15 ml/calf/day.

Sheep and Goats:

5 to 10 ml/adult animal/day; and
1 to 5 ml/young
animal/day.

Horses:

30 to 50 ml/adult animal/day;
and 5 to 15 ml/foal/day.

Generally, administer 1 ml/10 kg
b.w./day, for 3 days.

Shake well before use. Do not
administer more than 20 ml in
cattle or more than 10 ml in swine
and more than 5 ml in calves per
injection site to favour absorption
and dispersion.

WITHDRAWAL TIME

- For meat : 30 days.
- For milk : 2 days.

PACKAGING

Neutral glass vials of 50-mL and
100-mL, closed by a rubber
stopper and sealed with an
aluminum cover. Secondary
package: carton box.



BRIT PHARMA LTD

71-75 Shelton Street, Covent
Garden, London, WC2H 9JQ
T: +44 161 818 8992
E: info@brit-pharma.co.uk
W: www.brit-pharma.co.uk

STORE BELOW

30°C

STRICTLY FOR ANIMAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
STORE IN A DRY AND DARK PLACE.
KEEP BOTTLE CLOSED UPON USAGE.