



Buloxy 20 LA

Oxytetracycline Injection 20%

★★★★★ 5549 views

Content: 100ml:20g

Packing: 100ml

Application:Treatment of diseases caused by Oxtetracycline-susceptible organisms in cattle, sheep goats. Diseases including pneumonia shipping fever complex associated with Pasteur ella spp Haemophilus spp, etc.

Description

Composition

Each ml contains:
Oxytetracycline.....200mg

Indication

Treatment of diseases caused by oxytetracycline-susceptible organisms in cattle, sheep and goats. Diseases including pneumonia and shipping fever complex associated with Pasteurella spp and Haemophilus spp, infections bovine keratoconjunctivitis (pinkeye) bacterial enteritis(scours) caused by escherichia coli, wooden tongue caused by actinobacillus lignierisil; leptospirosis caused by leptospira pomoma; wound injections and acute metritis caused by stains of staphylococci and streptococci organisms sensitive to oxytetracycline.
Please see the enclosed that insert for more details.

Dosage and Administration

For intramuscular injection. The dose of the active principle Oxytetracycline amounts to 20 mg/kg bodyweight of the animal. The administration is carried out in the following volume:
Cattle,sheep and goats: 1ml / 10 kg bodyweight
If necessary, reinject with an interval of 3-5 days from the first administration.

Side Effects and Contraindications

Oxytetracycline injection is not intended for cats, dogs and horses. It should not be given to animals in late pregnancy, animals with severe damages of liver and kidneys and to animals oversensitive to Oxytetracycline. Sometimes a temporary swelling on the injection site occurs.

Withdrawal Time

Meat: 28 days
Milk: 7 days

Storage and Expired Time

Store in a dry and dark place, below 30°C
2 years

Packing