

FOR THE USE ONLY OF A VETERINARIAN OR
A HOSPITAL OR A LABORATORY OR A FARM

234307 R1

Rx

**Freeze Dried
Pellet of Natural
Glycoprotein
Human Chorionic
Gonadotrophin**



CHORULON®

NOT FOR HUMAN USE
FOR ANIMAL TREATMENT ONLY



SCHEDULE H PRESCRIPTION DRUG – CAUTION Not to be sold by retail without
the prescription of a Registered Veterinary Practitioner only

COMPOSITION

Each 1500 IU vial contains:
Gonadotrophin chorionic (hCG) 1500 I.U.
Mannitol 5.0 mg
Disodium phosphate dihydrate 0.315 mg
Sodium dihydrogen phosphate dihydrate 0.33mg

Solvent for reconstitution:
Disodium phosphate dihydrate 0.63 mg
Sodium dihydrogen phosphate dihydrate 0.66 mg
Water for injection, bulk 998.71 mg

PHARMACEUTICAL FORM

Powder for injectable solution, supplied with reconstitution solvent

TARGET SPECIES

Cattle (cows and heifers), horses (mares)

INDICATIONS

Induction of ovulation in mares.
Optimisation of fertility in cows and heifers (repeat breeders).
Treatment of follicular cysts in cows and heifers.

DOSAGE AND ADMINISTRATION

Animal species	Indication	Dosage and administration
Cow, heifer	improvement of conception rate	1500 IU, i.m. or i.v. at A.I. or mating and / or 4 to 6 days later
	Treatment of follicular cysts	3000 IU i.v.
Mare	induction of ovulation (follicles > 3,5 cm diameter)	1500 - 3000 IU i.m. or i.v. 24 hours before A.I. or mating

METHODS AND ROUTES OF ADMINISTRATION

Inject intramuscularly or intravenously

ADVICE ON CORRECT ADMINISTRATION

Chorulon can be used in early pregnant animals (upto 2 weeks following mating / insemination)

Do not use in animals in late pregnancy

Chorulon may be used during lactation

CONTRAINDICATIONS

None known

ADVERSE REACTIONS

In rare cases, as with all protein preparations, anaphylactic reactions may occur shortly after injection. When symptoms appear, adrenaline injection (1:1000) given intravenously or intramuscularly is the standard treatment. The administration of corticosteroid may also be indicated.

SPECIAL WARNINGS

Aseptic precautions should be adopted at the time of injection.

hCG has been used in human medicine for decades. There are no fundamental differences in effects of hCG in humans and animals: hCG can influence fertility in humans after injection. Care must be taken when using Chorulon to avoid self-injection.

In case of accidental self-injection, seek medical advice immediately and show the package insert or the label to the physician.

Accidental spillage on the skin should be washed-off immediately with soap and water.

WITHDRAWAL PERIODS

For milk and edible tissues: zero days

STORAGE CONDITIONS

Store at 2 to 8 °C

SHELF LIFE

36 months

After reconstitution, the product may be stored at +2°C to +8°C for upto 24 hours.

Manufactured by:

Intervet International B.V, Wim de Korverstraat 35 5831AN, Boxmeer, The Netherlands.

Imported and Marketed by:

Intervet India Pvt. Ltd.,

Building No. B-1, Godown No.12 to 16,

Ground Floor & Godown No. 12-A to 17-A First Floor, House No. 233,

Sagar Complex, Owali, Bhiwandi-14, Thane-

421302, Maharashtra, India.



PSC020502_02_R1

90 x 160 mm two sides printed

Black