



# CORTIVET DEPOT ACTION®

STEROID ANTI-INFLAMMATORY

|                                   | FORMULA |                         | PRESENTATION |
|-----------------------------------|---------|-------------------------|--------------|
| Dexamethasone 21 sodium phosphate | 25 mg   | Formulation agents q.s. | 10 ml        |
| Dexamethasone acetate             | 25 mg   |                         |              |

GLASS VIAL  
10 ml.  
100 ml.  
Sterile solution



## CORTIVET DEPOT ACTION®

**Cortivet Depot Action® is a long-acting anti-inflammatory** composed of two different dexamethasone salts: dexamethasone phosphate and dexamethasone acetate.

While dexamethasone phosphate is a rapidly released and metabolized corticosteroid, dexamethasone acetate is a depot glucocorticoid, with its release occurring between 2 and 14 days after application. The combination of these compounds has virtually no mineralocorticoid activity, making this product a very useful therapeutic tool for both acute and chronic inflammatory processes.

## INDICATIONS FOR USE

It is indicated for shock, stress, Neonatal Dysfunction Syndrome, traumatic brain injury, spinal cord compression, bronchospasm, allergic reactions, autoimmune syndromes, dermatitis and inflammation of the musculoskeletal system, and arthritis of traumatic origin via intra-articular injection.

## VIA OF ADMINISTRATION AND DOSAGE

**Dosage:** Horses of all ages: 1 to 4 ml per 100 kg body weight (0.05 to 0.2 mg/kg body weight).

**Frequency of Administration:** The treatment is a single application. Repeat administration is at the discretion of the attending veterinarian.

For intra-articular applications, the dosage will depend on the desired effect and the capacity of the joint being treated.

**Route of Administration:** Cortivet Depot Action® is a sterile injectable suspension and can be administered intramuscularly, subcutaneously, intra-articularly, epidurally, or by local infiltration.

## CONTRAINDICATIONS

Do not administer to immunosuppressed animals, concurrently with any vaccine, or in cases of fungal, viral, or bacterial infections where appropriate antibiotic therapy has not been initiated.

Do not use when gastric ulcer is suspected.

Do not administer intrasynovially if sepsis is observed in or around the joint, if there is a possibility of bone fracture or damaged cartilage, or if degenerative bone lesions or osteochondrosis are present.

## WARNINGS

Store the product between 15 and 25 °C.  
Keep out of reach of children and pets.

**For veterinary use only.**