



FARANTEL® EQUINOS 1000 KG

INTERNAL ANTIPARASITE
FOR EQUINES - ORAL USE

	FORMULA		PRESENTATION
Fenbendazole	15,0 g	DL – Metionine	12,5 g
Triclorfon	50 g	Excipients c.s.p.	100 g

SYRINGE

38 g. 

FARANTEL® EQUINOS 1000 KG

It is a vermicide, larvicide and ovicide in gastrointestinal and pulmonary parasitosis.

Due its ovicide effect it prevents contamination from pastures and stables, which is a key factor in getting rid of the infection.

Fenbendazole is a benzimidazole methylcarbonate of great anthelmintic capacity. Its action mechanism is through union to the tubulin protein and in this way it inhibits the polymerization of same to form microtubules. These form part of the cytoskeleton and, besides, participate in the transport of nutrients and the elimination of metabolites.

Trichlorfon is an organophosphate compound which confers a larvicide and ovicide function to the antiparasite formula by generating the irreversible inhibition of the enzyme acetylcholinesterase which leads to the blockade of the hydrolysis of acetylcholine in sites of cholinergic transmission. Due to the hepatic metabolism which this antiparasite has, a liver protector, Methionine is included. This works by providing methyl and sulphur to form choline and by eliminating certain toxic compounds in the organism.

INDICATIONS FOR USE

This antiparasite acts on nematodes, cestodes and trematodes in general, but it is excellent for combating parasitosis caused by *Ascaris*, *Parascaris*, *Strongylus vulgaris* and *edentatus*, *Strongyloides westeri*, *Habronema* and *Tridontotus*.

VIA OF ADMINISTRATION AND DOSAGE

- ½ syringe equivalent to 19 g of paste for a 500 kgv animal, equivalent to a dosage of each active ingredient:
- Fenbendazole: 5.7 mg/kgv
- Triclorfon: 19 mg/kgv
- DL-Methionine: 4.7 mg/kgv

CONTRAINDICATIONS

We recommend that it should not be used with female horses in the last three months of pregnancy nor in colts less than 6 months old.

Do not administer with other organophosphates or inhibitors of acetylcholinesterase.

Do not administer together with phenothazine tranquilizers nor with anesthetics 10 days before and after application.

WARNINGS

Store the product between 15 and 25 °C.
Keep out of reach of children and pets.

For veterinary use only.