



FENBENDAZOLE CALASTREMÉ

BROAD-SPECTRUM INTERNAL ANTIPARASITIC FOR ORAL USE, 10% SUSPENSION

Presentation: 1000 y 5000 mL drum

Formula: Every 100ml. contains: Fenbendazole 10 gr.; Vehicle q.s. 100 ml.

Indications:

Broad spectrum anthelmintic for ruminants, horses and pigs.

It is effective as an ovicidal and controls mature and immature stages of gastrointestinal and pulmonary helminths (Strongyloides, Pinworms, Ascarids, etc.)

Caution:

Treated animals should not be slaughtered earlier than 8 days after the Fenbendazole administration.

Restrictions:

Do not use it in females during the first days of pregnancy. Do not use it in dairy cattle during reproduction time.

Doses and routes of administration:

- In Cattle: 10 ml per 200 kg of live weight. In case of Dictyocaulosis and Ostertagia type II administer 15 ml per 200 kg of live weight. For both types of parasitosis, administer orally as a single dose.
- In Equines: 7.5 ml per 100 kg of live weight, orally as a single dose.
- In Sheep and Goats: 1 ml per 10 kg of live weight.
- In pigs: Porcine ascariasis: 1 ml per 10 kg of live weight, orally as a single dose.

NOTE: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.

Used containers must be discarded in accordance with current local legislation.

Store sheltered from light between 0° and 20°C.

Keep out of the reach of children and pets.

In the event of accidental poisoning attend the nearest poison control center.

