



Interflox-100

Solution for parenteral administration



Composition:

Contains per ml:

Enrofloxacin..... 100 mg.
Excipients ad 1 ml.

Description:

Enrofloxacin belongs to the group of quinolones and acts bactericidal against mainly Gram-negative bacteria like *Campylobacter*, *E. coli*, *Haemophilus*, *Pasteurella*, *Mycoplasma* and *Salmonella* spp.

Indications:

Gastrointestinal and respiratory infections caused by enrofloxacin sensitive micro-organisms, like *Campylobacter*, *E. coli*, *Haemophilus*, *Mycoplasma*, *Pasteurella* and *Salmonella* spp. in calves, cattle, sheep, goats and swine.

Contra-indications:

Hypersensitivity to enrofloxacin.

Administration to animals with a seriously impaired hepatic and/or renal function.

Concurrent administration of tetracyclines, chloramphenicol, macrolides and lincosamides.

Side effects:

Administration to young animals during growth can cause cartilage lesions in joints.

Hypersensitivity reactions may occur.

Dosage:

For intramuscular or subcutaneous administration:

Calves, cattle, sheep and goats : 1 ml per 20 - 40 kg body weight for 3 - 5 days
Swine : 1 ml per 20 - 40 kg body weight for 3 - 5 days.

Withdrawal times:

- For meat:
Calves, cattle, sheep and goats : 21 days.
Swine : 14 days.
- For milk : 4 days.

Packaging:

Vial of 50 and 100 ml.

