



Interspectin-L

Solution for parenteral administration

Composition:

Contains per ml:

Spectinomycin base	100 mg.
Lincomycin base.....	50 mg.
Excipients ad	1 ml.

Description:

The combination of lincomycin and spectinomycin acts additive and in some cases synergistic. Spectinomycin acts bacteriostatic or bactericidal, depending on the dose, against mainly Gram-negative bacteria like *Campylobacter*, *E. coli* and *Salmonella* spp. Lincomycin acts bacteriostatic against mainly Gram-positive bacteria like *Mycoplasma*, *Treponema*, *Staphylococcus* and *Streptococcus* spp. Cross-resistance of lincomycin with macrolides can occur.

Indications:

Gastrointestinal and respiratory infections caused by lincomycin and spectinomycin sensitive micro-organisms, like *Campylobacter*, *E. coli*, *Mycoplasma*, *Salmonella*, *Staphylococcus*, *Streptococcus* and *Treponema* spp. in calves, cats, dogs, goats, sheep and swine.

Contra-indications:

Hypersensitivity to lincomycin and/or spectinomycin.

Administration to animals with an impaired renal and/or hepatic function.

Concurrent administration of penicillins, cephalosporins, quinolones and cycloserine.

Side effects:

Hypersensitivity reactions.

Shortly after injection a slight pain, itching or diarrhoea can occur.

Dosage:

For intramuscular administration:

Calves	:	1 ml per 10 kg body weight for 4 days.
Goats and sheep	:	1 ml per 10 kg body weight for 3 days.
Swine	:	1 ml per 10 kg body weight for 3 - 7 days.
Cats and dogs	:	1 ml per 5 kg body weight for 3 - 5 days, with a maximum of 21 days.

Withdrawal times:

- For meat:

Calves, goats, sheep and swine : 14 days.

- For milk : 3 days.

Packaging:

Vial of 50 ml and 100 ml.

