



Net Content: 250 ml

KAOFIN

REGISTRATION NUMBER Q-7972-100

Gentamicin, Sulfamethoxazole, Trimetoprim and Loperamide

Broad spectrum antimicrobial and
antidiarrheal combination

Oral Suspension

FORMULA:

Every ml contains:

Gentamicin.....	12 mg
Sulfamethoxazole.....	100 mg
Trimethoprim.....	20 mg
Loperamide	0.5 mg
Excipient (Kaolin, Pectin) c.b.p.....	1 ml

CONSULT WITH YOUR VETERINARIAN

MEDICAL PRESCRIPTION IS NECESSARY FOR SELL

INDICATIONS:

Antidiarrheal and broad spectrum antimicrobial useful in infections caused by Gram positive and Gram negative microorganisms such as; *Escherichia coli*, *Pasteurella* spp, *Klebsiella* spp, *Pseudomonas* spp, *Proteus* spp, *Campylobacter* spp, *Vibrio* spp, *Treponemaspp* and *Clostridium*spp, as well as in some protozoan infections: *Trichomonas*, *Isospora* and *Eimeria*, micoplasmas and rickettsias sensitive to the action of the formula. Specifically used as an adjuvant for the treatment of various diarrheas such as: mechanical viral and bacterial, acute and in cases where it is desired to reduce intestinal motility.

SPECIES:

Cattle, Sheep, Goats, Swines, Canines and Felines.

DOSES:

Cattle (milk and meat) Sheep, Goats (milk and Meat), Pigs and Canines:
1 ml per 5 kg of body weight every 12 hours (equivalent to Gentamicin 2.4 mg /kg of body weight, Sulfamethoxazole 20 mg/kg of body weight, Trimethoprim 4 mg/kg of body weight and Loperamide 0.1 mg /kg of body weight).

Felines: 0.5 ml per 5 kg of body weight every 12 hours (equivalent to Gentamicin 1.2 mg/kg of body weight, Sulfamethoxazole 10 mg/kg of body weight, Trimethoprim 2 mg/kg of body weight and Loperamide 0.05 mg/kg of body weight).

ROUTE OF ADMINISTRATION:

Oral

WITHDRAWAL PERIOD:

Cattle, sheep and goats 10 days

Swines and piglets 10 days

MILK: Cattle, sheep and goats 7 days

WARNINGS AND/OR RESTRICTIONS:

All opioids such as loperamide should be used with caution in patients with hypothyroidism, renal failure, Addison's Syndrome, geriatric or frail patients, with glaucoma, or with secondary diarrhea caused by toxins and/or Salmonella spp., intestinal obstruction, or abdominal trauma. They are also contraindicated in cases where the patient is hypersensitive to narcotic analgesics or who are being treated with amino oxidase inhibitors. It should be used with caution in patients with head injuries or increased pressure, respiratory diseases and/or liver problems.

Keep up in a dry, cool place, in a range of 15-30°C and protected from light direct solar. No left within the reach of children and domestic animals. All unused veterinary medicine, waste derived from it and empty containers must be disposed of safely and in accordance with local regulations, avoiding contaminating water sources. **Product for exclusive use in Veterinary Medicine.**

VETERINARY USE

MADE IN MEXICO BY:

PET'S PHARMA DE MÉXICO S.A. DE C.V.

AV. Sor Juana Inés de la Cruz n° 580

Col. Benito Juárez, Nezahualcóyotl,

Estado de México P.C. 57000

Telephone: 55 57 43 38 39

ventas@petspharma.com.mx

petspharma.com.mx