



Net Content: 60 and 100 ml

KAOFIN

APPROVAL NUMBER Q-7972-100

Gentamicin, Sulfamethoxazole, Trimetoprim and Loperamide

Wide spectrum antimicrobial combination
and antidiarrheal

Oral Suspension

FORMULA:

Every ml contains:

Gentamicin.....	12 mg
Sulfamethoxazole.....	100 mg
Trimethoprim.....	20 mg
Loperamide	0.5 mg
Excipient (Kaolin, Pectin) c.b.p.....	1 ml

CONSULT WITH YOUR VETERINARIAN

**MEDICAL PRESCRIPTION IS
NECESSARY FOR SELL**

INDICATIONS:

Antidiarrheal and broad spectrum antimicrobial useful in infections caused by Gram positive, Gram negative microorganisms such as; *Escherichia coli*, *Pasteurella* spp, *Klebsiella* spp, *Pseudomonas* spp, *Proteus* spp, *Campylobacter* spp, *Vibrio* spp, *Treponema* spp and *Clostridium* spp, as well as in some protozoan infections: *Trichomonas*, *Isospora* and *Eimeria*, mycoplasmas and rickettsias sensitive to the action of the formula. It is specifically used as a coadjuvant for the treatment of diverse diarrheas such as: viral and bacterial mechanical diarrheas, acute diarrheas and in cases where it is desired to decrease the intestinal motility of the same.

SPECIES:

Bovine, Sheep, goats, pigs, canine and feline.

DOSES:

Cattle (milk and meat), sheep, goats (milk and meat), swine and canines:
1 ml per 5 kg body weight every 12 hours (equivalent to Gentamicin 2.4 mg /kg body weight, Sulfamethoxazole 20 mg /kg body weight, Trimethoprim 4 mg /kg body weight and Loperamide 0.1 mg /kg body weight).

Felines:

0.5 ml per 5kg body weight every 12 hours (equivalent to Gentamicin 1.2 mg /kg body weight, Sulfamethoxazole 10 mg /kg body weight, Trimethoprim 2 mg /kg body weight and Loperamide 0.05 mg /kg body weight).

ROUTE OF ADMINISTRATION:

Oral

WITHDRAWAL PERIOD:

CARNE:

Bovinos, Ovinos y Caprinos 10 días

Cerdos y lechones 10 días

LECHE:

Bovinos, Ovinos y Caprinos 7 días

WARNINGS AND/OR RESTRICTIONS:

All opioids such as loperamide should be used with caution in patients with hypothyroidism, renal insufficiency, Addison's Syndrome, geriatric or frail, with glaucoma, or with secondary diarrhea caused by toxins and/or Salmonella spp, intestinal obstruction, or abdominal trauma. They are also contraindicated in cases where the patient is hypersensitive to narcotic analgesics or is being treated with amino oxidase inhibitors. It should be used with caution in patients with head injuries or increased pressure, respiratory diseases and/or with hepatic problems. Keep in a dry, cool place at no more than 30°C and protected from direct sunlight. Keep out of reach of children, domestic animals and handicapped persons. **Product for exclusive use in Veterinary Medicine.**

VETERINARY USE

MADE IN MEXICO BY:

PET'S PHARMA DE MÉXICO S.A. DE C.V.

AV. Sor Juana Inés de la Cruz n° 580

Col. Benito Juárez, Nezahualcóyotl,

Estado de México P.C. 57000

Telephone: 55 57 43 38 39

ventas@petspharma.com.mx

petspharma.com.mx