



Net Content: 20 tablets

MICROFLOX 2.5%

APPROVAL NUMBER Q-7972-111

SOLUGEL **Enrofloxacin**

Wide Spectrum Antimicrobial
Oral Solution

FORMULA:

Every ml contains:

Enrofloxacin 25 mg

Excipient c.b.p. 1 tablet

CONSULT WITH YOUR VETERINARIAN

MEDICAL PRESCRIPTION IS NECESSARY FOR SELL

DESCRIPTION:

MICROFLOX 2.5% solugel contains micronized enrofloxacin, which allows better digestive absorption, bioavailability, faster tissue penetration and better liposolubility. It acts against Gram positive bacteria, Gram negative Mycoplasma spp, and some Rickettsiae. Inhibits cell division and bacterial growth rate, prevents tissue colonization and promotes phagocytosis.

INDICATIONS:

It is recommended for otitis externa, septicemia, enterobacteria, prevention and treatment of osteomyelitis, skin, urogenital, respiratory, prostatic and wound infections. It has post-antibiotic effect on Staphylococcus intermedius, Enterobacter spp, Psuedomonas aeruginosa, Klebsiella pneumoniae, Pasteurella multocida, Escherichia coli, Salmonella typhimurium, Streptococcus aureus and Rickettsia rickettsi.

SPECIES:

Canines, Felines, Mustelids (Ferrets), Small Mammals (rabbits,

hedgehogs, chinchillas, gerbils, guinea pigs, hamsters, mice and rats). Ornamental birds (passerines, psittacines and pigeons), Swine, bovine meat producers. NOT indicated for birds producing eggs for human consumption.

DOSES.

Swine (piglets): 5 mg/kg body weight, 1mL/5 kg body weight, Beef cattle: 2.5 mg/kg body weight, 1mL/10 kg body weight, every 24 hours for 3 to 5 days.

Canines and Felines: 5 mg/kg body weight, 1mL/5 kg body weight, every 24 hours for 3-5 days.

Mustelids (Ferrets) and Small Mammals: (rabbits, hedgehogs, chinchillas, gerbils, guinea pigs, hamsters, mice and rats): 5-10 mg/kg body weight, which is equivalent to 1mL/2.5-5 kg body weight, every 12 hours for 14 days.

Ornamental birds (passerines, psittacines and pigeons) 5-10 mg/kg body weight, equivalent to 1mL/2.5-5 kg body weight, every 8 hours for 3 to 5 days.

WITHDRAWAL PERIOD:

Swine 10 days.

Beef cattle 5 days.

ROUTE OF ADMINISTRATION:

Oral.

WARNINGS AND/OR RESTRICTIONS:

Any Enrofloxacin produces erosion of articular cartilage, so its use is not recommended in growing animals. Do not exceed the recommended dose because it decreases its effectiveness and increases genotoxicity on eukaryotic cells. DO NOT use in dairy cattle destined for human consumption. DO NOT administer in poultry producing eggs for human consumption, nor in replacement birds 14 days before the beginning of laying. Keep in a dry, cool place at no more than 30°C and protected from direct sunlight. Keep out of reach of children, domestic animals and handicapped persons. **Product for exclusive use in Veterinary Medicine.**

VETERINARY USE

MADE IN MEXICO BY:

PET'S PHARMA DE MÉXICO S.A. DE C.V.

AV. Sor Juana Inés de la Cruz n° 580

Col. Benito Juárez, Nezahualcóyotl,

Estado de México P.C. 57000

Telephone: 55 57 43 38 39

ventas@petspharma.com.mx

petspharma.com.mx