



NEUROLIS PLUS

INJECTABLE FOR SPORT HORSES | LOCAL ANESTHETIC

Presentation: 50 mL bottle

Composition: Each 100 cc contains: Ammonium Chloride 2.00%; Benzyl Alcohol 0.75%; Bidistilled water q.s. 100 ml.

Used for the treatment of sciatic pain, intercostal pain, neuritis and pain of nervous origin in different places of equines' limbs.

Indications:

It is indicated to relieve muscle pain. Indicated in cases of myositis, myalgia, muscle contractures. Its effect lasts 2 to 3 weeks and begins to be noticeable 2 hours after injection. It is indicated in large muscle groups such as the shoulder, rump or back. It can be used pure or together with antibiotics and local anesthetics.

Local effects: The horse may show discomfort after the infiltration. It may be associated with lidocaine to avoid symptoms.

Caution: It must be used exclusively by a veterinarian.

Contraindications: Injuries which may worsen with the blockage: fissures, fractures, tendinitis or infectious processes. Small inflammatory reactions may occur at the place of injection, which disappear shortly after the application.

Routes of administration: Exclusively intramuscular (infiltrations)

Paravertebral: Cervical 2-3 ml; Dorsal: 5-10 ml; Lumbar 5-10 ml; Sacral: 3-5 ml.

Caudal: 10 ml/ Sciatic: 10 ml/ Local: infiltrative 5-10 ml.

One application every 30 days.

DO NOT USE IN EQUINES INTENDED FOR HUMAN CONSUMPTION.

Keep out of the reach of children and pets.

In the event of accidental poisoning attend the nearest poison control center.

Storage: between 4° and 30°C.

Used containers must be discarded in accordance with current local legislation.



NEUROLIS PLUS

Injectable neurolytic
for sport horses

50 mL
VETERINARY USE

To be sold with a
veterinary prescription



Calastremé
PRODUCTOS VETERINARIOS

COMPOSITION: each 100 cc contains:
Ammonium Chloride 2.00%; Benzyl Alcohol
0.75%; Bidistilled water q.s. 100 ml.

ROUTES OF ADMINISTRATION AND
PARAVERTEBRAL DOSAGE: Exclusively
intramuscular (infiltrations)

Cervical 2-3 ml.
Dorsal 5-10 ml.
Lumbar 5-10 ml.
Sacral 3-5 ml.
Caudal 10 ml.
Sciatic 10 ml
Local (infiltrative) 5-10 ml.

One application every 30 days. DO NOT USE
IN EQUINES INTENDED FOR HUMAN
CONSUMPTION.

NOTE: The distributor is not responsible for the
consequences arising from improper use of
the product, since it will be used in
circumstances beyond the distributor's control.
Keep out of reach of children and pets.
In the event of accidental poisoning attend the
nearest poison control center.

Storage: between 4° and 30°C.

Indications:

It is intended for the relief of muscle pain.

Indicated in cases of myositis, myalgia,
muscle contractures.

Produced:

Expires:



SENSA Cert. N° 05-260
Prod. establishment N° 8533



ARGENTINE PRODUCT



PROTECT
THE ENVIRONMENT

Manufactured & Distributed by:

Establecimiento Calastremé S.A.

La Paz 2150 B1640CYB Mtz., Bs. As.,

Argentina. Tel.: (011) 4580-9999

Technical Director: Dr. Carlos N. Nuñez

Vet. physician. Prof. registration N°: 1350

e-mail: info@calastreme.com.ar

www.calastreme.com.ar