

**WIDE SPECTRUM
LONG-ACTION ANTIBIOTIC
ANTI-INFLAMMATORY**

FORMULA

Each 100 ml contains:

Oxytetracycline dihydrate (*) . . . 32.4 g

Flunixin meglumine (**) 3.3 g

Excipients q.s.ad 100.0 ml

(*) equivalent to 30 g Oxytetracycline

(**) equivalent to 2 g Flunixin base



**Oximic
30% FM**

**OXITETRACYCLINE 30%
+ FLUNIXIN BASE 2%**

Veterinary Use

INDICATIONS

Oximic 30% FM is a long-acting antibiotic and anti-inflammatory. Synergic association of a wide spectrum and long-acting antibiotic (Oxytetracycline L.A.: up to 7 days in bovines, camelids and South American auenids) with a non-steroid analgesic and antipyretic anti-inflammatory (Flunixin), with anti-endotoxic effect for 24 to 36 hours, to optimize treatment in one single application.

Oxytetracycline, in all the species indicated: to promote a lasting and sustained action for the control and treatment of infectious conditions of the respiratory (bovine respiratory disease) and urinary systems and of local infections caused by Gram positive and Gram negative germs, such as: *Bordetella bronchiseptica*; *Actinomyces pyogenes*; *Erysipelothrix rhusopathiae*; *Bacillus anthracis*; *Clostridium spp.*; *Pasteurella spp.*; *Campylobacter spp.*; *Haemophilus spp.*; *Klebsiella spp.*; *Actinobacillus spp.*; *Actinomyces bovis*, *Leptospira pomona*; *Moraxella bovis*; *Erysipelotrix insidiosa*; *Corynebacterium spp.*; *Listeria spp.*; *Streptococcus spp.*; *Staphylococcus spp.*; *Anaplasma spp.*; *Dermatophilus congolensis* and certain Mycoplasmas, Rickettsias, Chlamydiae, and some pathogenic fungi and protozoa, sensitive to oxytetracycline.

Flunixin, in all the indicated species: for infections when an anti-inflammatory, antipyretic and anti-endotoxic effect is required. Treatment for inflammation of locomotive apparatus and musculoskeletal system. It is also used in inflammatory conditions associated to the "Bovine Respiratory Complex" due to its antipyretic action. Neonatal diarrhea, downer cow syndrome and for treatment of acute mastitis.

TARGET SPECIES

Bovines, South American camelids and auenids.

ROUTE OF ADMINISTRATION

It is administered through deep intramuscular parenteral route, preferably in the neck area (neck side area). Do not administer more than 10 ml per inoculation site.

DOSAGE

Suggested posology: 10 ml per 100 kg l.w. every 5 to 7 days (equivalent to 30 mg/kg of l.w. oxytetracycline and to 2 mg/kg of l.w. flunixin).

Oxytetracycline, in all the indicated species:

Dose: 20 mg oxytetracycline/kg. l.w. every 2 to 4 days (equivalent to 10 ml per 150 Kg l.w.) for light infections;

30 mg/kg l.w. every 5 to 7 days (equivalent to 10 ml every 100 kg. l.w.) for moderate to severe infections.

Flunixin: 1.1 - 2.2 mg/kg. Administer every 5 to 7 days, the interval may vary according to the needs in each case and/or at veterinarian's discretion.

WITHDRAWAL PERIOD

Do not administer in dairy females whose milk is intended for human consumption.

Do not slaughter animals whose meat is intended for human consumption until the following time periods have elapsed:

- When administering a dose of 20 mg/kg. l.w. = until 28 days after last treatment.
- When administering a dose of 30 mg/kg. l.w. = until 35 days after last treatment.

PRECAUTIONS

Oxytetracycline: prolonged use thereof may induce the development of mycosis due to the suppression of the normal flora. In gestating females it may induce an abnormal color in the teeth of new-borns. Due to an inotropic and negative vasodilation action, it may result in lowering blood pressure.

Flunixin, intra-arterial injection may cause adverse reactions. They are brief and disappear without treatment. Occasionally, intramuscular administration in bovines may cause irritation at inoculation site.

Do not administer in gestating females.

Use in animals younger than 6 weeks old is not recommended, as it may cause adverse reactions.

Do not use in severe cases of renal, hepatic or cardiac injury, or in animals with parasitic injuries of the gastrointestinal mucous membrane.

Do not use in the presence of colics caused by the ileum and associated to dehydration.

Use of product concomitantly with other non-steroid anti-inflammatory products is not recommended.

Keep out of the reach of children and pets. Never re-use empty containers.

Once used, containers and wastes should be disposed of and destroyed adequately, either by burning them in safe places or burying them far away from lakes, streams, water courses, etc.

Store in a dry, cool place; away from sunlight and preferably between 5 and 25° C.

VETERINARY USE

MADE IN URUGUAY

M.G.A.P. (Ministry of Livestock, Agriculture and Fishing of Uruguay) reg. no.: A-4587

Owner and manufacturer

LABORATORIOS MICROSULES URUGUAY S.A.

Ruta 101 km.28 - Cno. al Paso Escobar s/n

Pueblo Cap. J. A. Artigas - Canelones - Uruguay - C.P. 91.001

Tel. (00598-2) 2886761* - Fax (00598-2) 2886760

Technician in charge: Dr. Pablo R. Bringas

www.laboratoriosmicrosules.com

