

POLYEQUAN inj. ad us. vet. Serum against foal paralysis

1. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT
Bioveta, a.s., Komenského 212,
683 23 Ivanovice na Hané, Czech Republic

2. NAME OF THE VETERINARY MEDICINAL PRODUCT

POLYEQUAN inj. ad us. vet.
Serum against foal paralysis

3. STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENT(S)

Composition – 100 ml: Immunoserum anti-
Escherichia coli, *Salmonella abortus equi*,
Streptococcus equi, *Actinobacillus equinum*
nativum 100 ml, Phenolum.

4. INDICATION

Passive immunization of foals when breeding is endangered or in case of illness induced by *E. coli*, *Salmonella abortus equi*, *Streptococcus equi*, *Actinobacillus equinum nativum* germs.

5. CONTRAINDICATIONS

Not known.

6. ADVERSE REACTIONS

Not known.

7. TARGET SPECIES

Foals.

8. DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION

Dosage:

Preventive: Foals aged up to 1 week 25 ml
Foals aged 1 week and above 50 ml
Therapeutic: double doses should be applied, half of it subcutaneously and the rest intravenously.

The application can be repeated, if necessary. Subcutaneous or intravenous application should be used.

9. ADVICE ON CORRECT ADMINISTRATION

Shake the contents of the vial well before use.

10. WITHDRAWAL PERIOD

Without withdrawal period.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children.
Keep in a dry and dark place at of 2 to 8 °C.

12. SPECIAL WARNING(S)

Are not.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

14. DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED
August 2006

15. OTHER INFORMATION

The antibodies contained in equine serum ensure the immunity in animals for 10 – 14 days. Dynamics of resorption and usability of antibodies during various application methods is particularly important.

The antibodies applied intravenously are contained directly in blood serum and their usability is 90%.

The serum against the foal paralysis applied subcutaneously is absorbed only slowly but usability of globulins is 10% only.

Interaction with other medicinal products and other forms of interaction

None known.

Inkompatibility

None known.

Package: 50ml, 100 ml.

Shelf Life: 2 years.

Shelf-life after first opening – 10 hours

For animal treatment only - to be supplied only on veterinary prescription.



Dimensions (mm):

Colours:

Prepared by:

Date:

A6 (105 × 148 mm) duplex printing
black

Hanáková

15. 3. 2019

Approved by:

Date: