

SERGON 500 IU/ml**powder for preparation of injection solution with solvent****1. NAME AND ADDRESS OF THE MARKETING****AUTHORISATION HOLDER AND OF THE MANUFACTURING AUTHORISATION HOLDER RESPONSIBLE FOR BATCH RELEASE, IF DIFFERENT**

Marketing authorisation holder and manufacturer:

Bioveta, a. s.,
Komenského 212,
683 23 Ivanovice na Hané,
Czech Republic**2. NAME OF THE VETERINARY MEDICINAL PRODUCT**SERGON 500 IU/ml powder for preparation of injection solution with solvent
Gonadotrophinum sericum equinum**3. STATEMENT OF THE ACTIVE SUBSTANCES AND OTHER INGREDIENTS**

1 ml of reconstituted preparation contains:

Active substances:

Gonadotrophinum sericum equinum – 500 IU

Excipients:Methylparaben (E 218) – 1.8 mg
Propylparaben (E 216) – 0.2 mg**4. INDICATIONS**

Anestrus, induction and synchronization of heat.

5. CONTRAINDICATIONS

None.

6. ADVERSE REACTIONS

As the preparation is purified, anaphylactoid reactions are minimized.

7. TARGET SPECIES

Cow, heifers, sows, sheep, goat, bitch, rabbit.

8. DOSAGE FOR EACH SPECIES, ROUTE AND METHOD OF ADMINISTRATION

1) cows and heifers:

1000 – 3000 IU

2) sows: 500 – 1000 IU

Application schedule:

sows – induction of sexual cycle and increasing of the number of piglets per litter (1st – 2nd day after estrum)– quiet heat (10th day after piglet weaning)

Young sows – anestrus, quiet heat (8 – 10 months old)

– heat induction (aged 6 months or 90 kg live weight. Insemination in the first heat after treatment.

Insemination in the second heat results in more numerous litters.)

3) sheep and goats:

500 IU (suitable immediately after removing of intra-vaginal tampons)

4) bitches: 250 – 500 IU

5) rabbit: 25 – 50 IU (insemination 3rd and 5th day after application)

Administration: intramuscular or subcutaneous.

9. ADVICE ON CORRECT ADMINISTRATION

Lyophilisate dissolve by added Dissolvent A.

10. WITHDRAWAL PERIOD

None.

11. SPECIAL STORAGE PRECAUTIONS

Keep out of the reach and sight of children.

Keep in refrigerator (2 – 8°C).

Protect from light.

Do not use after the expiry date stated on the label.

Shelf life after dilution or re-constitution as instructed: 24 hours.

12. SPECIAL WARNING

If ingested accidentally, do not induce vomiting. Wash the mouth with large amounts of water.

If the person is conscious, give him/her enough water to drink.

In case of undesired effects, seek medical aid.

In case of skin contact, wash with water and soap. In case of skin inflammation, seek medical aid.

In case of eye contact, wash with sufficient amounts of water, including under the lids, for at least 15 minutes. In case of inflammation, seek medical aid.

If injected incidentally, seek medical aid immediately and show the physician leaflet info or label.

Use suitable protective clothes to prevent incidental application.

13. SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS, IF ANY

Any unused veterinary medicinal preparation or waste originated from this preparation must be disposed of according to local legal regulations.

14. DATE OF REVISION OF THE TEXT

2016

15. OTHER INFORMATIONFor animal treatment only!
Veterinary medicinal product subject to prescription!

Dimensions (mm): A6 (105 × 148 mm) duplex printing
 Colours: black
 Prepared by: Vlková
 Date: 30. 9. 2016

Approved by:

Date: