

ADVERSE REACTION:

When used according to manufacturer recommendations, the product does not cause any adverse reaction.

WITHDRAWAL PERIOD: This product does not require a withdrawal period.

PRESENTATION:

Amber glass vial containing 10 mL display with 25 amber glass vials containing 10 mL each.

Transport the product at a temperature no more than 25°C. Store the product protected from sunlight at temperatures between 15°C and 30°C, and keep away from children and pets.

SALE AND USE UNDER VETERINARIAN PRESCRIPTION

Registered at the Ministry of Agriculture, Livestock, and Food Supply under Number SP 000211-9.000002 in 03/23/2017

Technical Responsible:

Dr. Jose Antonio Dell'Aqua Junior

Veterinarian

CRMV/SP 11220

OWNED AND MANUFACTURED BY:

BOTUPHARMA INDUSTRIA E COMERCIO DE PRODUTOS VETERINARIOS LTDA.

Coronel Fonseca, 2111 – Jardim Bom Pastor

Botucatu, SÃO PAULO 18603-495

CNPJ: 11.195.056/0001-65

Ins. Estadual: 224.197.376.119

Phone: +55 (14) 3112-9900

Fax: +55 (14) 3112-9904

www.botupharma.com.br

vendas@botupharma.com.br

MADE IN BRAZIL

BATCH NUMBER AND MANUFACTURING DATE:

See carton and label.

EXPIRATION DATE:

02 (two) years after the manufacturing date.

Botupharma
INDUSTRIA E COMERCIO DE PRODUTOS VETERINARIOS

Strelin
Injectable Histrelin

Strelin

Injectable Histrelin

**VETERINARY USE****FORMULA:**

Each 1 mL contains:

Histrelin Acetate250 µg

Vehicle q.s.....1 mL

INDICATION:

Strelin is a potent synthetic GnRH analogue (Gonadotropin Release Hormone). It is indicated for mares to induce ovulation and synchronization.

ROUTE OF ADMINISTRATION:

Injectable product exclusively for deep intramuscular injection.

HOW TO USE AND DOSAGE:

Strelin should be administered to mares that have ovarian follicles equal to or greater than 33 mm in diameter, determined by transrectal palpation and ultrasonography, in a single dose of 1 mL (250 µg). During the starting and ending periods of the breeding season, the dose may be doubled (2 mL to 500 µg) at the veterinarian's discretion, for greater product efficiency.

INTERVAL BETWEEN DOSES AND TREATMENT DURATION:

There is no strictly defined interval for product reapplication. Usually, its administration is done in a single dose. Reapplication is subject to the veterinarian's decision, who is responsible for diagnosis, prescription and follow-up of the animal in treatment.

CONTRAINDICATION AND LIMITATION OF USE:

Do not use in animals that are hypersensitive to any of product components. Repeated application in short time intervals may cause inhibition of gonadotrophins by hypophysis. Use before expiration date.

CAUTION AND PRECAUTION IN USE:

The product should be used according to the manufacturer's recommendation. Always use sterile needles and syringes, complying with good aseptic practices.