



SULFALKYL INYECTABLE

TREATMENT FOR INFLAMMATION

Presentación: 50 and 100 mL bottle

Formula: DMSO 100%

TARGET SPECIES: SPORTING EQUINES

Indications:

Arthritis, osteoarthritis, tendinitis, bursitis, synovitis, inflammatory processes in general. Inflammation of the CNS due to spinal cord pressure caused by trauma, or thoracic-lumbar myalgia of a neurological origin, pleuritis, septic arthritis, encephalomyelitis.

Dosage and method of administration:

Slow IV exclusively. Apply 20% diluted in physiological solution, at a rate of 0.1 to 0.25 g/kg of weight, depending on the desired effect and the criteria of the intervening Veterinary Doctor.
Apply every 12 hours. Treatment may be extended up to 7 days.

Contraindications:

Avoid using it together with organophosphates due to its anticholinergic action. Do not administer to pregnant females. Do not administer to animals intended for human consumption.

WARNING:

Product may crystallize at low temperatures. In such a case, warm the closed bottle on a double boiler (BM) until the crystallization disappears. It may cause local hypersensitivity.

Keep out of the reach of children and pets.

Store between 15° and 25°C and sheltered from light.

Used containers must be discarded according to current local legislation. Discard leftover product once the bottle has been opened.

In the event of accidental poisoning attend the nearest poison control center.

NOTE: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.



Calastremé

SULFALKYL INJECTABLE

100 mL

VETERINARY USE

To be sold with a
veterinary prescription



Calastremé
PRODUCTOS VETERINARIOS

COMPOSITION: DMSO... 100%

TARGET SPECIES: EQUINES

INDICATIONS: Arthritis, osteoarthritis, tendinitis, bursitis, synovitis, inflammatory processes in general. Inflammation of the CNS due to spinal cord pressure caused by trauma, or thoracic-lumbar myalgia of a neurological origin, pleuritis, septic arthritis, encephalomyelitis.

DOSAGE AND METHOD OF ADMINISTRATION: Slow IV exclusively. Apply 20% diluted in physiological solution, at a rate of 0.1 to 0.25 g/kg of weight, depending on the desired effect and the criteria of the intervening Veterinary Doctor. Apply every 12 hours. Treatment may be extended up to 7 days.

CONTRAINDICATIONS: Avoid using it together with organophosphates due to its anticholinergic action. Do not administer to pregnant females. Do not administer to animals intended for human consumption.

WARNING: Product may crystallize at low temperatures. In such a case, warm the closed bottle on a double boiler (BM) until the crystallization disappears. It may cause local hypersensitivity.

Keep out of the reach of children and pets. Store between 15° and 25°C and sheltered from light. Used containers must be discarded according to current local legislation. Discard leftover product once the bottle has been opened. In the event of accidental poisoning attend the nearest poison control center.

NOTE: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.

Series:

Produced:

Expires:



SENASA Cert. Nº 11-206

Producing establishment Nº 8533



PROTECT
THE ENVIRONMENT



ARGENTINE PRODUCT

Manufactured & Distributed by:
Establecimiento Calastremé S.A.

La Paz 2150 B1640CYB Mtz., Bs. As., Argentina.

Tel.: (011) 4580-9999

Technical Director: Dr. Carlos N. Nuñez. Vet. physician. Prof. registration Nº: 1350

e-mail: info@calastreme.com.ar | www.calastreme.com.ar