



TIAMINA HAMPTON

VITAMINI

Presentation: 100 mL vial bottle

Composition: Thiamine hydrochloride 30 g.; Excipients q.s. 100 ml.

Description:

Thiamine vitamin supplement. This vitamin's contribution improves the mechanisms of complete oxidation of glucose, especially in those tissues energetically dependent on glucose or isotate-pyruvate (heart and nervous system), which under physiological demands (pregnancy, lactation, fever, energetic exercise or breeding demands) increase its need.

Indications:

To prevent or treat functional disorders connected to the deficit or increased demand for thiamine (neuritis, polyneuritis, sciatic syndromes, rheumatism, pain and to reinforce glycogen fixation)

Route of administration: Slow intravenous, with the animal at rest.

Dose:

Preventive: 6 to 10 ml every 48 hours for 15 days

Curative: 10 ml daily for 15 days.

Species: Sport equines. Do not administer to animals intended for human consumption.

Warning and caution:

The intravenous administration must be carried out slowly, since anaphylactic reactions have been exceptionally described. Protect from light and excessive heat.

NOTE: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.

Keep out of the reach of children and pets.

In the event of accidental poisoning attend the nearest poison control center.

Storage: Between 5° and 25° C and sheltered from light.

Used containers must be discarded in accordance with current local legislation.

Discard leftover product once the container has been opened.



TIAMINA HAMPTON

VITAMIN
Injectable solution

100 mL

VETERINARY USE
To be sold with a
veterinary prescription



Calastremé
PRODUCTOS VETERINARIOS

COMPOSITION:

Thiamine Hydrochloride 30 g.
Excipients q.s. 100 ml

PRESENTATION: 100 ml vial

DESCRIPTION: Thiamine vitamin supplement. This vitamin's contribution improves the mechanisms of complete oxidation of glucose, especially in those tissues energetically dependent on glucose or isotate-pyruvate (heart and nervous system), which under physiological demands (pregnancy, lactation, fever, energetic exercise or breeding demands) increase its need.

INDICATIONS: To prevent or treat functional disorders connected to the deficit or increased demand for thiamine (neuritis, polyneuritis, sciatic syndromes, rheumatism, pain and to reinforce glycogen fixation).

ROUTE OF ADMINISTRATION: Slow intravenous, with the animal at rest.

DOSE: Preventive: 6 to 10 ml every 48 hours for 15 days
Curative: 10 ml daily for 15 days.

SPECIES: Sport equines. Do not administer to animals intended for human consumption.

WARNING AND CAUTION: The intravenous administration must be carried out slowly, since anaphylactic reactions have been exceptionally described. Protect from light and excessive heat.

NOTE: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.

**Keep out of the reach of children and pets
In the event of accidental poisoning attend the nearest
poison control center.**

Storage: Between 5° and 25° C and sheltered from light.
Used containers must be discarded in accordance with current local legislation. Discard leftover product once the container has been opened.

Series:

Produced:

Expires:



SENASA Cert. N° 79.478



Producing establishment N° 8533



ARGENTINE PRODUCT



PROTECT
THE ENVIRONMENT

Manufactured & Distributed by:
Establecimiento Calastremé S.A.

La Paz 2150 B1640CYB Mtz., Bs. As., Argentina.

Tel.: (011) 4580-9999

Tech. Dir.: Dr. Carlos N. Nuñez. Vet. Ph. Prof. registr. N°: 1350

e-mail: info@calastreme.com.ar | www.calastreme.com.ar