

ORAL POWDER BRONCHODILATOR

Equines not intended for human consumption.

Formula:

Clenbuterol hydrochloride 0,002 g.

Excipients q.s. 100,000 g.

Indications for use: Ventical is used as a bronchodilator for the treatment of respiratory disease in horses. It relieves Emphysema, bronchospasms, accumulation of mucus, chronic obstructive pulmonary disease (COPD), bronchitis.

Dosage and Method of administration: Administer a measure (10 g) per 200 kg bodyweight orally every 12 hours for 14 days unless otherwise stipulated by a veterinarian physician.

Precautions: Do not use simultaneously with corticosteroids.

Contraindications:

Do not use in horses intended for human consumption.

Do not use in pregnant or lactating mares

Note: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.

Keep out of the reach of children and pets.

Store between 15° and 25°C and sheltered from light.

Used containers must be discarded in accordance with current local legislation.

In the event of accidental poisoning attend the nearest poison control center.

 SENASA Cert. N° 05.042

 Prod. establ. N°: 8533

ARGENTINE PRODUCT 

Produced and distributed by Establecimiento Calastremé S.A.

La PAZ 2150-(B1640CYB) Martínez, Bs. As. - Arg. Ph: 011 4580-9999

Technical Director: Dr. Carlos N. Nuñez. Vet. Doc. - Med. Reg. P.B.A 1350

e-mail: info@calastreme.com.ar | www.calastreme.com.ar

VENTICAL

Clenbuterol

To be sold with a filed
veterinary prescription

VETERINARY USE

500 g

MADE IN ARGENTINA



Calastremé
PRODUCTOS VETERINARIOS



Sport Horses

ANTICATARRHAL - BRONCHODILATORS

VENTICAL

CLENBUTEROL | ORAL POWDER BRONCHODILATOR

Presentation: 500 g Pot

Formula: Clenbuterol hydrochloride 0.002 g; Excipients q.s. 100.000 g.

Indications for use:

Ventical is used as a bronchodilator for the treatment of respiratory disease in horses. It relieves Emphysema, bronchospasms, accumulation of mucus, chronic obstructive pulmonary disease (COPD), bronchitis.

Dosage and method of administration:

Administer a measure (10 g) per 200 kg bodyweight orally every 12 hours for 14 days unless otherwise stipulated by a veterinarian physician.

Precautions:

Do not use simultaneously with corticosteroids.

Contraindications:

Do not use in horses intended for human consumption.

Do not use in pregnant or lactating mares.

NOTE: The distributor is not responsible for the consequences arising from improper use of the product, since it will be used in circumstances beyond the distributor's control.

Keep out of the reach of children and pets.

Store between 15° and 25°C and sheltered from light.

Used containers must be discarded in accordance with current local legislation.

In the event of accidental poisoning attend the nearest poison control center.



Calastremé