

VITAPLASTIN Forte

Mineral food complement for cattle, sheep, goats, pigs, dogs, silver foxes, rabbits, poultry and exotic birds

Manufacturer: Bioveta, a.s., Komenského
212, 683 23 Ivanovice na Hané, Czech
Republic
production site 1, Tyršova 409/40,
Ivanovice na Hané 683 23, Czech Republic

**Approval identification number of the
production site:** a CZ 800585-01

Composition: Calcium carbonate, calcium
dihydrogen and hydrogen phosphate
monohydrate (monocalcium phosphate);
calcium magnesium carbonate

Analytical constituents:

Humidity:	10.0 %
Calcium (Ca):	300.0 g/kg
Phosphorus (P):	76.0 g/kg
Magnesium (Mg):	7.0 g/kg
Sodium (Na):	1 g/kg

Supplemental substances:

Nutritional supplemental substances

(compounds of trace elements):

iron (3b103 – iron (II) sulphate
monohydrate ($\text{FeSO}_4 \cdot \text{H}_2\text{O}$), 3b106 – iron (II)
chelate of amino acids hydrate
($\text{Fe}(\text{x})_{1-3} \cdot \text{n H}_2\text{O}$)) 8000 mg/kg
iodine 3b201 (potassium iodide (KI))
35 mg/kg
copper (3b404 – copper (II) oxide (CuO),
3b406 – copper (II) chelate of amino acids
hydrate ($\text{Cu}(\text{x})_{1-3} \cdot \text{n H}_2\text{O}$)) 280 mg/kg
manganese (3b502 – manganese (II) oxide
(MnO), 3b504 – manganese chelate of
amino acids hydrate ($\text{Mn}(\text{x})_{1-3} \cdot \text{n H}_2\text{O}$)) 225
mg/kg
Zinc (3b603 – Zinc oxide (ZnO), 3b607 –
zinc chelate of glycine, hydrate (solid) (Zn
(x) $_{1-3} \cdot \text{n H}_2\text{O}$)) 1800 mg/kg

Senzorical complementary substances:

2a Other colourants
E 172 Iron oxide (Fe_2O_3)
2b Natural products – botanically defined
Vanilla aroma 0.6 g/1 kg

Indication: To supplement the minerals in
the feed ration.

Administration: Oral, in feed.

Dosage:

Cattle: 30-60 g per day, i.e. one heaped
tablespoon 1-2 times daily. In large-
capacity breeding farms one 1/2 litre bottle
of powder 1-2 times daily per each 10-14
cows.

Calves: 10-30 g per day, i.e. a heaped
teaspoon 1-3 times daily, depending on the
size.

Sheep, goats: 10 g per day, i.e. a flat
teaspoon 2 times daily.

Lambs: 1-3 times a day one knife point to
the smallest ones; 3-5 g per day, i.e. less
than or a flat teaspoon once daily to older
ones.

Pigs: one flat teaspoon 1-2 times daily, or
one heaped teaspoon per each 3 heads 3
times daily /5-10 g/.

Piglets: 1-3 times daily one knife point to
the smallest ones; 3-5 g per day, i.e. less
than or a flat teaspoon once daily, or one
heaped teaspoon per each 3 heads 1-2
times daily to older ones.

Dogs, silver foxes: 1-5 g per day, i.e. one
knife point 2 times daily, up to one flat
teaspoon once daily.

Rabbits: one knife point once daily.

Poultry: 0.5-1.5 g per day, i.e. one knife
point 1-3 times daily, or one flat to heaped
teaspoon per 10 hens or 50-100 chickens
once daily. Proportionately more to
ducklings or goslings, depending on their
body weight.

Exotic birds: one knife point into 100 g of
soft feed.

Withdrawal periods: Not applicable.

Pack size: 1 kg, 5 kg

For animal treatment only!

bioveta